

Article

Dilatatsion kardiomiopatiya bilan og‘rigan bemorlarda kasallik kechishining og‘irlik darajasi va tizimli yallig‘lanish o‘rtasidagi bog‘liqlik

T.A. Abdullayev ¹ , M.M. Rahmatov ¹ , X.A. G‘ulomov ^{*1} 

¹ Yurak yetishmovchiligi va miokardning nokoronarogen kasalliklari bo‘limi, Respublika ixtisoslashtirilgan kardiologiya ilmiy-amaliy tibbiyot markazi, Toshkent, 100052, O‘zbekiston

escardio@mail.ru (T.A.), mhmdjnrhmtv@gmail.com (M.R.), Hgulamov93.08@gmail.com (X.G.)

* Correspondence: mhmdjnrhmtv@gmail.com; I.: +998901223639 (M.R.)

Annotatsiya:

Maqsad. Dilatatsion kardiomiopatiya bilan og‘rigan bemorlarda klinik-gemodinamik ko‘rsatkichlar va yallig‘lanish markerlari (C-reaktiv oqsil, IL-6, O‘NO- α) o‘rtasidagi bog‘liqlikni o‘rganish.

Materiallar va usullar. Tadqiqotga dilatatsion kardiomiopatiya bilan og‘rigan hamda SYuYe NYHA bo‘yicha II-IV FS bilan asoratlangan 42 nafar ($38,5 \pm 1,2$ yil) bemor kiritildi. Bemorlar ikki guruhga ajratildi: I guruh – FS II ($n=12$) va II guruh – FS III-IV ($n=30$).

Natijalar. SYuYe og‘irlik darajasi ortishi bilan (FS III-IV) bemorlar qon zardobida C-reaktiv oqsil ($7,6 \pm 1,1$ mg/l va $4,1 \pm 0,6$ mg/l, $p<0,05$), IL-6 ($7,5 \pm 0,8$ pg/ml va $4,6 \pm 0,2$ pg/ml, $p<0,01$) va O‘NO- α ($8,1 \pm 0,9$ pg/ml va $5,6 \pm 0,7$ pg/ml, $p<0,05$) miqdori oshgan, bu esa chap qorincha dilatatsiyasi (YaSO⁺ 58 ± 1 , $p<0,05$) va otish fraksiyasining pasayishi ($34,7 \pm 1,8\%$, $p<0,05$) bilan bog‘liq bo‘lgan.

Xulosa. Yallig‘lanish markerlari DKMP og‘irligi va prognozini baholashda muhim diagnostik ahamiyatga ega.

Kalit so‘zlar: Dilatatsion kardiomiopatiya, surunkali yurak yetishmovchiligi, yallig‘lanish, C-reaktiv oqsil, interleykin-6, o‘sma nekrozi omili- α , kolxitsin.

Association between disease severity and systemic inflammation in patients with dilated cardiomyopathy

Timur A. Abdullayev ¹ , Mahmud M. Rakhmatov ¹ , Khumoyun A. Gulomov ^{*1} 

¹ Department of Heart Failure and Non-Coronary Myocardial Diseases, Republican Specialized Scientific and Practical Medical Center of Cardiology, Tashkent, 100052, Uzbekistan

escardio@mail.ru (T.A.), mhmdjnrhmtv@gmail.com (M.R.), Hgulamov93.08@gmail.com (Kh.G.)

Abstract:

Aim. The aim of this study was to investigate the relationship between clinical-hemodynamic parameters and inflammatory markers (C-reactive protein, interleukin-6, tumor necrosis factor- α) in patients with dilated cardiomyopathy.

Materials and methods. The study included 42 patients (mean age 38.5 ± 1.2 years) with dilated cardiomyopathy and chronic heart failure classified as NYHA functional class II–IV. The patients were divided into two groups: Group I – class II ($n=12$) and Group II – class III–IV ($n=30$).

Results. As heart failure severity increased (class III–IV), serum levels of C-reactive protein (7.6 ± 1.1 mg/L vs. 4.1 ± 0.6 mg/L, $p<0.05$), IL-6 (7.5 ± 0.8 pg/mL vs. 4.6 ± 0.2 pg/mL, $p<0.01$), and TNF- α (8.1 ± 0.9 pg/mL vs. 5.6 ± 0.7 pg/mL, $p<0.05$) were significantly elevated. These changes were associated with left ventricular dilatation (LVEDD 58 ± 1 mm, $p<0.05$) and reduced ejection fraction ($34.7 \pm 1.8\%$, $p<0.05$).

Iqtibos: T.A. Abdullayev, M.M. Rahmatov, X.A. G‘ulomov. Dilatatsion

kardiomiopatiya bilan og‘rigan bemorlarda kasallik kechishining og‘irlik darajasi va tizimli yallig‘lanish o‘rtasidagi bog‘liqlik. 2025, 2,4, 6.

<https://doi.org/10.70626/cardiouz-2025-2-00065>

Olingan: 10.10.2025

Tuzatilgan: 18.10.2025

Qabul qilingan: 25.11.2025

Nashr qilingan: 10.12.2025

Copyright: © 2025 by the authors. Submitted to for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Conclusion. The findings suggest that inflammatory markers play a crucial role in assessing the severity and prognosis of dilated cardiomyopathy and chronic heart failure.

Keywords: Dilated cardiomyopathy, chronic heart failure, inflammation, C-reactive protein, interleukin-6, tumor necrosis factor- α , colchicine.

Kirish

2023-yilda Yevropa Kardiologiya Jamiyati (ESC) tomonidan ishlab chiqilgan kardiomiopatiyalarni davolash bo'yicha yangi klinik tavsiyalar so'nggi 15 yil mobaynida tibbiyotning turli sohalari hamkorligida olib borilgan jadal ilmiy faoliyat natijasida yaratilgan bo'lib, kardiologik jamiyatning klinik amaliyotiga joriy qilinmoqda [1]. Ushbu xujjat ESC tomonidan kardiomiopatiyalarni tasniflashning yangi bayonotini e'lon qilish bilan boshlangan bo'lib, unga shuningdek genetik maslahat va genetik testlash, kardiomiopatiyalarni aniqlash bo'yicha klinik-diagnostik yondashuvlar, gipertrofik kardiomiopatiya bo'yicha klinik tavsiyalar, gipokinetik nodilatatsion kardiomiopatiya bo'yicha takliflar, yurak amiloidozi bo'yicha bayonot hamda genetik variantlarni interpretatsiya qilish va ularni klinik amaliyotda qo'llash imkoniyatlarini qamrab olgan muhim hujjatlar hamrohlik qildi [2,8]. Ishlab chiqilgan yangi tasnifga ko'ra kardiomiopatiyalar 5 ta fenotipga ajratilgan — gipertrofik kardiomiopatiya (GKMP), dilatatsion kardiomiopatiya (DKMP), restriktiv kardiomiopatiya (RKMP), aritmogen o'ng qorincha kardiomiopatiyasi (AO'QKMP) va chap qorinchaning nodilatatsion kardiomiopatiyasi (ChQNDKMP).

Kardiomiopatiya fenotiplari orasida dilatatsion kardiomiopatiya chap qorincha otish fraksiyasi pasaygan yurak yetishmovchiligi (ChQOFpasayganYuYe) rivojlanishiga olib keluvchi eng keng tarqalgan sabablaridan biri sifatida tan olingan bo'lib, butun dunyoda yurak transplantatsiyasi bajarilishiga asosiy ko'rsatma hisoblanadi [9]. Dilatatsion kardiomiopatiya — asosan chap qorinchaning (ChQ) dilatatsiyasi hamda uning global yoki regional sistolik disfunktsiyaning mavjudligi bilan tavsiflanib, ushbu holatni faqatgina yuklama sharoitlarining buzilishi (masalan, gipertoniya, yurak qopqoqlari kasalliklari, tug'ma yurak nuqsonlari) yoki yurak ishemik kasalligi bilan izohlab bo'lmaydi [10]. So'ngi ma'lumotlarga ko'ra DKMP ning tarqalish darajasi har 100 000 aholiga 36 holatni tashkil etadi, yillik kasallanish esa har 100 000 aholiga 7 ta holat deb baholanadi [9,11]. Kasallik prognozi uning kelib chiqish sababi, klinik bosqichi va qo'llanilgan davolash choralarining samaradorligiga bog'liq holda jiddiy farq qiladi. Standart terapiya sharoitida 5 yillik yashab qolish ko'rsatkichi taxminan 50–60% atrofida bo'lsa-da, refrakter yurak yetishmovchiligi yoki qorinchalar aritmiyasi mavjud bemorlarda o'lim xavfi ortadi [12].

Surunkali yurak yetishmovchiligini (SYuYe) davolashning zamonaviy strategiyalari medikamentoz terapiya, elektr qurilmalar (CRT-D, implantatsiya qilinadigan kardioverter-defibrillyator) va ChQ yordamchi qurilmalarini o'rnatish yoki yurak transplantatsiyani o'z ichiga oladi [13]. Hozirgi kunda surunkali yurak yetishmovchiligiga olib kelgan etiologik omillardan qat'iy nazar ChQ OF pasaygan YuYe ni medikamentoz davolashda to'rt komponentli standart terapiya qo'llanilmoqda. Bular angiotenzin-aylantiruvchi fermenti ingibitorlari (iAAF)/angiotenzin retseptor va neprilizin ingibitorlari (ARNI), beta-blokatorlar (BB), mineralokortikoid retseptorlari antagonistlari (MRA), 2-tip natriy-glyukoza kotransporter ingibitorlari (SGLT-2), shuningdek shish sindromi bo'lganda qovuzloq diuretiklaridir [13]. Biroq optimal medikamentoz terapiya qo'llanilishiga qaramasdan DKMP bilan og'rigan bemorlarning hayot prognozi xali ham noxushligicha saqlanib qolmoqda, bu esa kasallik patogenezi yanada chuqurroq o'rganishni hamda yangi davolash usullarini ishlab chiqishni talab etadi.

So'nggi yillarda SYuYe rivojlanishining yangi kontsepsiyasi taklif etilgan bo'lib, unga ko'ra tizimli yallig'lanish yurak yetishmovchiligining noxush kechishi va yuqori yurak-qon tomir xavfining prediktori sifatida qaralmoqda [14]. SYuYe natijasida yuzaga keladigan mikrotsirkulyatsiyadagi jiddiy buzilishlar to'qimalardagi makrofaglar va monotsitlarning nospetsifik faollashuviga, ular esa o'z navbatida yallig'lanishdagi sitokinlar — o'sma nekrozi omili-alfa (O'NO- α), interleykin-1 (IL-1), interleykin-6 (IL-6) va boshqalar sintezining kuchayishiga olib keladi. Ushbu sitokinlar ChQ disfunktsiyaning rivojlanishiga ta'sir etuvchi muhim patogenetik mexanizmlardan biri sifatida qaralmoqda [15].

Maqsad

Dilatatsion kardiomiopatiya bilan ogʻrigan bemorlarda klinik-gemodinamik koʻrsatkichlar va qon zardobidagi yalligʻlanish markyori – C-reaktiv oqsil (CRO) va yalligʻlanisholdi sitokinlari (IL-6, OʻNO- α) miqdori oʻrtasidagi bogʻliqlikni oʻrganish.

Materiallar va usullar

Tadqiqot Xelsinki deklaratsiyasi tamoyillariga muvofiq ravishda amalga oshirildi. Protokol RIKIATM huzuridagi Etik qoʻmita tomonidan maʼqullandi. Tadqiqotga jalb etilishidan oldin barcha ishtirokchilardan yozma ravishda rozilik olindi. Tadqiqotga RIKIATM 7-Miokardning nokoronarogen kasalliklari va pulmonologiya boʻlimidagi DKMP tashxisi qoʻyilgan hamda SYuYe NYHA boʻyicha II-IV FS bilan asoratlangan 42 nafar (25 erkak va 17 ayol) 18 yoshdan 57 yoshgacha boʻlgan (oʻrtacha yosh $38,5 \pm 1,2$ yil) bemorlar olindi. Kardiomiopatiyalarni davolash boʻyicha ESC 2023 tavsiyalariga muvofiq bemorlarga DKMP tashxisi qoʻyildi. Barcha bemorlardan anamnestik maʼlumotlar yigʻildi, KHBSH shkalasi yordamida klinik holat baholandi, 6 daqiqalik yurish testi (6DYuT) oʻtkazildi va uning asosida NYHA tasnifiga koʻra yurak yetishmovchiligi funktsional sinfi aniqlandi. Barcha bemorlar SYuYe funktsional sinfiga koʻra ikki guruhga boʻlindi: I guruh – FS II boʻlgan bemorlar ($n=12$); II guruh – FS III-IV boʻlgan bemorlar ($n=30$). Shuningdek, barcha bemorlarda umumklinik tekshiruvlar: standart 12 tarmoqli EKG tekshiruvi, transtorakal ExoKG tekshiruvi bajarildi. Bundan tashqari barcha bemorlar qon zardobida yalligʻlanish markyori (CRO) va yalligʻlanisholdi sitokinlari (IL-6, OʻNO- α) miqdori aniqlandi. Tadqiqotga infeksiyon, allergik, onkologik va immun tizim kasalliklari, shuningdek, oʻpkalar surunkali obstruktiv kasalligi bilan ogʻrigan bemorlar qoʻshilmadi, negaki ushbu kasalliklarda yalligʻlanish markyorlari va sitokinlar miqdori yuqori boʻladi.

Olingan maʼlumotlarning statistik tahlili MedCalc dasturiy taʼminoti yordamida amalga oshirildi. Aniqlangan koʻrsatkichlar orasida arifmetik oʻrtacha qiymat (M), standart ogʻish (SD) va arifmetik oʻrtacha qiymatning standart xatosi (m) hisoblab chiqildi. Ahamiyatli farqlar Styudentning t-kriteriysi asosida aniqlandi. $p<0,05$ boʻlgan holatlar statistik jihatdan ishonchli deb hisoblandi. Maʼlumotlar $M \pm m$ koʻrinishida keltirildi.

Natijalar

Ikkala guruhdagi bemorlarning boshlangʻich klinik va demografik koʻrsatkichlari taqqoslanganda ular oʻrtasida ishinochli farqlar aniqlanmadi. I va II guruhdagi bemorlarning KHBSH shkalasi boʻyicha toʻplangan ballar yigʻindisi mos ravishda $7,3 \pm 0,7$ ball va $8,9 \pm 0,3$ ballni tashkil etdi. Shuningdek, 6DYuT davomida bosib oʻtilgan masofa I va II guruhlarda mos ravishda $350,4 \pm 19,7$ metr va $248,8 \pm 13,2$ metrni tashkil etdi (Jadval 1).

Table 1. Initial clinical and demographic parameters ($M \pm m$) in patients with DCM according to the severity of HF

Jadval 1. DKMP bilan ogʻrigan bemorlarda SYuYe ogʻirlik darajasiga koʻra boshlangʻich klinik va demografik koʻrsatkichlar ($M \pm m$)

Koʻrsatkich	I guruh FS II ($n=12$)	II guruh FS III-IV ($n=30$)	p
Yosh, yil	$41,4 \pm 2,6$	$41,5 \pm 2,1$	$p>0,05$
Kasallikning davomiyli, oy	$9,4 \pm 3,8$	$8,7 \pm 1,6$	$p>0,05$
YuQS, daqiqa	$87,2 \pm 3,3$	$93,7 \pm 2,7$	$p>0,05$
YuQS, daqiqa	$87,2 \pm 3,3$	$93,7 \pm 2,7$	$p>0,05$
SAB, mm.sim.ust.	$110,8 \pm 4,3$	$108,4 \pm 2,9$	$p>0,05$
DAB, mm.sim.ust.	$72,5 \pm 3,3$	$69,2 \pm 1,9$	$p>0,05$

Izoh: n – bemorlar soni; p - I guruhga nisbatan ishonchlik darajasi; KHBSH – klinik holatni baholash shkalasi; 6DYuT – 6 daqiqalik yurish testi; YuQS – yurak qisqarishlari soni; SAB – sistolik arterial bosim; DAB – diastolik arterial bosim.

Ikkala guruhdagi bemorlarning ExoKG tekshiruv natijalari taqqoslanganda quyidagilar aniqlandi (Jadval 2). II guruhdagi bemorlarning (FS III-IV) ExoKG koʻrsatkichlari I guruhdagi bemorlarniki (FS II) bilan taqqoslanganda deyarli barcha koʻrsatkichlarda yaqqol salbiy oʻzgarishlar aniqlandi, biroq faqatgina YaSOʻ, YaSH, OF va ChB oʻlchami koʻrsatkichlari orasidagi farq ishonchli deb baholandi ($p<0,05$). Jumladan, YaSOʻ koʻrsatkichlari I va II guruhlarda mos ravishda 52 ± 2 mm

va 58 ± 1 mm ni tashkil etdi; YaSH ko'rsatkichlari I va II guruhlarda mos ravishda $134,5 \pm 13,4$ ml va $170,1 \pm 10,7$ ml ni tashkil etdi; ChQ OF ko'rsatkichlari I va II guruhlarda mos ravishda $43,1 \pm 3,5$ % va $34,7 \pm 1,8$ % ni tashkil etdi; ChB o'lchami ko'rsatkichlari I va II guruhlarda mos ravishda 41 ± 2 mm va 54 ± 5 mm ni tashkil etdi. Qolgan ko'rsatkichlar o'rtasida ishonchli farqlar aniqlanmadi.

Table 2. Cardiac hemodynamic parameters ($M \pm m$) in patients with DCM according to the severity of HF
Jadval 2. DKMP bilan og'rikan bemorlarda SYuYe og'irlik darajasiga yurakichi gemodinamikasi ko'rsatkichlari ($M \pm m$)

Ko'rsatkich	I guruh FS II (n=12)	II guruh FS III-IV (n=30)	p
QAT, mm	$9,7 \pm 0,5$	$8,9 \pm 0,3$	$p > 0,05$
ChQOD, mm	$9,8 \pm 0,5$	$9,5 \pm 0,3$	$p > 0,05$
YaDO', mm	67 ± 2	70 ± 1	$p > 0,05$
YaSO', mm	52 ± 2	$58 \pm 1^*$	$p < 0,05$
YaDH, ml	$236,1 \pm 15,9$	$261,8 \pm 14,9$	$p > 0,05$
YaSH, ml	$134,5 \pm 13,4$	$170,1 \pm 10,7^*$	$p < 0,05$
OF, %	$43,1 \pm 3,5$	$34,7 \pm 1,8^*$	$p < 0,05$
ChB, mm	41 ± 2	$54 \pm 5^*$	$p < 0,05$
O'Q, mm	28 ± 3	35 ± 3	$p > 0,05$

Izoh: n – bemorlar soni; p - I guruhga nisbatan ishochlik darajasi; YaDO' – yakuniy diastolik o'lcham; YaSO' – yakuniy sistolik o'lcham; YaDH – yakuniy diastolik hajm; YaSH – yakuniy sistolik hajm; OF – otish fraktsiyasi; ChB – chap bo'lmaccha; O'Q – o'ng qorincha; QAT – qorinchalar aro to'siq; ChQOD – chap qorincha orqa devori.

Biz shuningdek bemorlar qon zardobidagi yallig'lanisholdi sitokinlari (IL-6, O'NO- α ,) va CRO miqdorini tekshirdik va ikkala guruh o'rtasida taqqoslama tahlil o'tkazildi. Bemorlarda yurak yetishmovchiligining og'irlik darajasi ortib borishi bilan yallig'lanish markyori va sitokinlar miqdori ham ishonchli darajada ko'tarilib borganligi aniqlandi. Xususan, I guruhdagi bemorlarda (FS II) CRO miqdori $4,1 \pm 0,6$ mg/l ni tashkil etgan bo'lsa, II (FS III-IV) guruhdagi bemorlar qon zardobidagi CRO miqdori $7,6 \pm 1,1$ mg/l ni tashkil etdi hamda ushbu ko'rsatkich II guruh bemorlarida I guruhdagi bemorlarnikiga nisbatan 85,4% ga yuqori bo'ldi ($p < 0,05$). Ikkala guruhdagi bemorlar qon zardobidagi IL-6 miqdorining taqqoslama tahlili o'tkazilganda, I guruhdagi bemorlar qon zardobida IL-6 miqdori $4,6 \pm 0,2$ pg/ml ni tashkil etdi, II guruhdagi bemorlarda esa $7,5 \pm 0,8$ pg/ml ni tashkil etdi hamda II guruh bemorlarida I guruhdagi bemorlarnikidan ishonchli ravishda 63 % ga yuqori ekanligi aniqlandi ($p < 0,01$). YuE funktsional sinfi ortib borishi bilan bemorlar qon zardobidagi O'NO- α miqdori ham ortib borgan hamda I va II guruhlarda mos ravishda $5,6 \pm 0,7$ pg/ml va $8,1 \pm 0,9$ pg/ml ni tashkil etdi. II guruhdagi bemorlar qon zardobidagi O'NO- α miqdori I guruhdagi bemorlarnikiga nisbatan 44,6% ga yuqori ekanligi aniqlandi ($p < 0,05$).

Table 3. Serum pro-inflammatory cytokines (IL-6, TNF- α) and CRP levels ($M \pm m$) in patients with DCM according to the severity of HF

Jadval 3. DKMP bilan og'rikan bemorlarda SYuYe og'irlik darajasiga ko'ra qon zardobidagi yallig'lanisholdi sitokinlari (IL-6, O'NO- α) va CRO miqdori ($M \pm m$)

Ko'rsatkich	I guruh FS II (n=12)	II guruh FS III-IV (n=30)	p
CRO, mg/l	$4,1 \pm 0,6$	$7,6 \pm 1,1^*$	$p < 0,05$
IL-6, pg/ml	$4,6 \pm 0,2$	$7,5 \pm 0,8^{**}$	$p < 0,01$
O'NO- α , pg/ml	$5,6 \pm 0,7$	$8,1 \pm 0,9^*$	$p < 0,05$
O'NO- $\alpha > 6,0$ pg/ml	5 (41,6%)	21 (70%)	

Izoh: n – bemorlar soni; p - I guruhga nisbatan ishochlik darajasi; CRO – C-reaktiv oqsil; O'NO- α – o'sma nekrozi omili- α ; IL-6 – interleykin-6.

Keltirilgan ma'lumotlar SYuYe og'irlik darajasi hamda qon zardobidagi C-reaktiv oqsil va IL-6, O'NO- α kabi yallig'lanisholdi sitokinlari o'rtasida yaqqol musbat bog'liqlik borligini ko'rsatadi.

Munozara

Etiologiyasidan qat'iy nazar, yurak yetishmovchiligining rivojlanishi ham mahalliy, ham tizimli yallig'lanish kaskadlarining faollashuvi bilan bog'liq [16]. Miokardit yoki yallig'lanishli kardiomiopatiya bilan og'rigan bemorlarda yallig'lanish yurak yetishmovchiligining asosiy sababi hisoblanadi [17]. Bundan tashqari, sitokinlar ta'sirida yuzaga kelgan yallig'lanish o'tkir stress-indutsirlangan "Takotsubo" kardiomiopatiyasi rivojlanishida ham ishtirok etishi aniqlangan [18]. Yurak shikastlanishiga javoban yallig'lanish reaksiyasi dastlab adaptiv xarakterga ega bo'ladi, biroq ortiqcha va uzoq davom etuvchi yallig'lanish javobi noadaptiv jarayonga aylanib, yurak yetishmovchiligi patogenezida hal qiluvchi omillardan biri bo'lib qoladi. Bir qator tadqiqot natijalariga ko'ra, yurak yetishmovchiligi mavjud bemorlar qon zardobida o'sma nekrozi omili- α ($O^*NO-\alpha$), interleykin-6 (IL-6), interleykin-8 (IL-8) kabi yallig'lanisholdi sitokinlari, hamda C-reaktiv oqsil (CRO) miqdori yuqori bo'lgan [19,20]. Uzoq muddatli subklinik tizimli yallig'lanish yurak mushagidagi yallig'lanish javobining shakllanishida hal qiluvchi o'rin egallashi aniqlangan [21]. Faollashgan to'qima makrofaglari ham yallig'lanisholdi sitokinlari va xemokinlarni sintezlab, miokardda yallig'lanish jarayonini kuchaytiradi hamda miokard fibrozlanishi va kardiomiopatiya rivojlanishi uchun sharoit yaratadi [22]. Jumladan, idiopatik DKMP rivojlangan bemorlarning deyarli yarmida biopsiya materiallarida yallig'lanisholdi sitokinlari miqdorining oshganligi va miokardda limfotsitar infiltratsiya aniqlangan [23].

DKMP bilan og'rigan bemorlar miokard to'qimasida doimiy ravishda $O^*NO-\alpha$ ekspressiyasining oshishi qayd etilgan [24]. Belenkov Yu.N. va hammualliflarning tadqiqotida kardiomiotsitlarning o'zi $O^*NO-\alpha$ ni ishlab chiqarish qobiliyatiga ega ekanligi va bunda sitokin miqdori to'g'ridan-to'g'ri miokardning zo'riqish darajasiga bog'liq ekanligi isbotlangan [25]. Yurak yetishmovchiligi bo'lgan bemorlar qon zardobida $O^*NO-\alpha$ ning yuqori darajasi ChQ OF pasaygan va ChQ OF saqlangan subpopulyatsiyalarda o'lim ko'rsatkichlarining oshishi bilan bog'liq bo'lgan [26].

Qon zardobidagi IL-1 miqdori turli etiologik omillarga (miokard infarkti, chap qorinchaga ortiqcha bosim yuklamasi va kardiomiopatiyalarga) bog'liq yurak yetishmovchiligining eksperimental modellarida doimiy ravishda oshganligi kuzatiladi [27–29]. Yurak yetishmovchiligida immun hujayralari, fibroblastlar, endoteliy hujayralari va kardiomiotsitlar IL-1 sintezi va faollashuviga hissa qo'shishi aniqlangan [30–32]. IL-1 tizimining faollashuvi turli mexanik yo'llar orqali yurakning sistolik va diastolik faoliyatining buzilishiga olib keladi hamda YuE patogenezida muhim ahamiyat kasb etishi mumkin [33]. Masalan, IL-1 siklik AMF ga bog'liq bo'lmagan mexanizm orqali L-tipdagi kaltsiy kanallarining beta-adrenergik reaksiyasini pasaytirishi isbotlangan [34]. Bundan tashqari, IL-1 kardiomiotsitlarda azot oksidi sintezlovchi ferment ekspressiyasini oshiradi, bu esa azot oksidi faolligining ortishiga va miokard qisqaruvchanlik qobiliyatining pasayishiga olib keladi [35]. 2017 yilda italiyalik olimlar tomonidan DKMP bilan og'rigan bemorlarda inflammasomalar va tizimli yallig'lanish biomarkerlari ahamiyatini o'rganishga qaratilgan tadqiqot o'tkazilgan. Bu tadqiqot doirasida 156 nafar ambulator bemor tekshirilgan (o'rtacha yosh — 58 yosh, 77% — erkaklar, ChQ OF medianasi — 35%, NUP medianasi — 189 pg/ml, IL-1 — 1,08 pg/ml, IL-6 — 1,7 pg/ml, IL-10 — 2,7 pg/ml). 89,6 oylik kuzatuv davomida 35 nafar bemor (22%) vafot etgan yoki yurak transplantatsiyasi amaliyoti bajarilgan bo'lib, ularning ko'pchiligida NYHA bo'yicha III FS, bo'lmachalar fibrillyatsiyasi (BF), ChQ OF ancha past va BNP kontsentratsiyasi ancha yuqori ekanligi qayd etilgan. IL-1, IL-6 va IL-10 miqdori yaxshi va yomon prognozli guruhlar o'rtasida sezilarli farq qilmagan. Shuningdek, IL-1 miqdori NYHA sinflari yoki ChQ OF kvartillari orasida ham ahamiyatli farq ko'rsatmagan. Biroq, ko'p omilli tahlilda IL-1 barcha sabablar natijasida yuzaga kelgan o'limning mustaqil va kuchli prediktori ekanligi aniqlangan [36].

Bir nechta klinik tadqiqotlar shuni ko'rsatdiki, yurak yetishmovchiligi bilan og'rigan bemorlarning miokard to'qimasida sog'lom odamlarnikiga nisbatan IL-6 ekspressiyasi yuqori bo'lgan [37]. Simptomsiz sistolik disfuntsiya bilan kechuvchi kasalliklarda IL-6 miqdorining ortishi aniqlangan, shuningdek, simptomsiz keksa bemorlar populyatsiyasida IL-6 va C-reaktiv oqsil (CRO) miqdorining ortishi yurak yetishmovchiligi holatlarini bashorat qilishda qo'llanilishi ko'rsatilgan [38,39]. Interleykin-6 hepatotsitlarda CRO ishlab chiqarilishining kuchli induktori bo'lib, u gipoksiyaga javoban bir necha turdagi hujayralar, shu jumladan, kardiomiotsitlar tomonidan ham ishlab chiqariladi [40]. TIME-CHF tadqiqoti turg'un surunkali yurak yetishmovchiligi bilan og'rigan

chap qorincha otish fraktsiyasi saqlangan va pasaygan bemorlar qon zardobida yuqori sezgirlikka ega C-reaktiv oqsil miqdorining oshganligi ko'rsatilgan [41].

Bizning tadqiqotimizda DKMP bilan og'rikan bemorlarning qon zardobida yallig'lanisholdi sitokinlari (IL-6, O'NO- α) va yallig'lanish markeri (CRO) miqdori ishonchli darajada oshgani aniqlandi. Tadqiqotimizda yurak yetishmovchiligining og'irlik darajasi ortishiga bog'liq ravishda yurakning patologik remodellanish darajasi hamda qon zardobidagi sitokinlar miqdori va C-reaktiv oqsil miqdori oshganligi ko'rsatildi.

Yurak yetishmovchiligi patogenezida yallig'lanisholdi sitokinlari muhim o'rin egallashi, tizimli yallig'lanishga qarshi yo'naltirilgan maxsus terapiya usullarini qo'llashni taqozo etadi. An'anaviy tarzda yallig'lanish kasalliklarini davolashda qo'llanilgan kolxitsin, yurak-qon tomir kasalliklarini davolashda o'zining xavfsizligini isbotlagan hamda hozirgi kunga kelib istiqbolli terapevtik dori vositasi sifatida o'rganilmoqda. Kolxitsin mikronaychalarni nishonga olib, ularning yig'ilishini va ma'lum yallig'lanish markerlarining ishlab chiqarilishini ingibirleydi, shuningdek neytrofillar faoliyatiga ta'sir ko'rsatib, aterosklerotik pilakchalar shakllanishiga to'sqinlik qilishi mumkin [42]. Shu sababli kolxitsin yurak ishemik kasalligi (YuIK), yurak ritmi buzilishlari (bo'lmachalar fibrillyatsiyasi), so'nggi yillarda esa yurak yetishmovchiligi va insult kabi holatlarni davolashdagi imkoniyatlari o'rganilmoqda.

2014 yilda Deftereos va hammualiflar tomonidan o'tkazilgan tadqiqotda va 2024 yildagi COLICA tadqiqotida yurak yetishmovchiligi bilan og'rikan bemorlarda kolxitsinni qo'llash imkoniyatlari o'rganilgan. Deftereos tadqiqotida kolxitsinning turg'un yurak yetishmovchiligi bo'lgan bemorlarning klinik holatiga ta'siri o'rganilgan [43]. Bu tadqiqotda kolxitsin terapiyasi bemorlar qon zardobidagi yallig'lanish markerlarini kamaytirgan bo'lsada, biroq bemorlarning NYHA bo'yicha funktsional sinfiga, o'lim ko'rsatkichiga yoki gospitalizatsiyalar soniga ijobiy ta'sir ko'rsatmagan. COLICA tadqiqotida o'tkir dekompensatsiyalangan yurak yetishmovchiligi bo'lgan bemorlarda kolxitsinning klinik samaradorligini baholangan [44]. Asosiy yakuniy nuqta — 8 haftadan so'ng NT-proBNP miqdorining pasayishi bo'lib, bu ko'rsatkich kolxitsin va platsebo guruhlar o'rtasida farq qilmagan. Kolxitsin qabul qilgan bemorlarda yallig'lanish markerlari, jumladan CRO va IL-6 miqdori kamaygan bo'lsada, ammo yurak yetishmovchiligining zo'rayishi yoki qayta dekompensatsiya holatlari bo'yicha guruhlar o'rtasida statistik farq aniqlanmagan.

Shunga qaramay yurak yetishmovchiligini davolashda kolxitsinning ta'sirini o'rganishga qaratilgan yangi tadqiqotlar olib borilmoqda. Xuan Sun va hammualiflarining 2022-yilda AHA jurnalida chop etilgan tadqiqotida eksperimental sichqonlar modelida doksorubitsin yordamida chaqirilgan DKMP ni davolashda kolxitsinning kuchli yallig'lanishga qarshi ta'siri ko'rsatilgan [45]. Ushbu ta'sir SIRT2 orqali amalga oshirilib, NLRP3 inflammasomasining faollashuvini susaytirish natijasida DKMP rivojlanishini sekinlashtirgan. Tadqiqotdan quyidagi natijalar olingan: - kolxitsin miokarddagi yallig'lanishni, neytrofillar infiltratsiyasini va yallig'lanish sitokinlari (IL-1, IL-6, O'NO- α) miqdorini kamaytirgan; - kolxitsin SIRT2 ni inaktivatsiya bo'lishdan himoya qilgan va shu orqali NLRP3 ni deatsetillab, uning faollashuvini cheklagan; - kolxitsin yurak faoliyatini yaxshilagan, miokard remodellanishining oldini olgan, vazn yo'qotish va doksorubitsin orqali chaqirilgan o'lim darajasini kamaytirgan. Tadqiqotchilarning xulosasiga ko'ra kolxitsin DKMP rivojlanishini sekinlashtirishda samarali natija ko'rsatgan bo'lib, uni yallig'lanish etiologiyali yurak yetishmovchiligini davolashda arzon, xavfsiz va istiqbolli vosita sifatida qo'llash imkoniyatini ilmiy asoslaydi.

Xulosa

Ko'plab ilmiy tadqiqot natijalari shundan dalolat beradiki, etiologiyadan qat'iy nazar, yurak yetishmovchiligining rivojlanishi sitokinlar va yallig'lanish markyorlari induktsiyasi bilan bog'liq bo'lib, ular yurakning salbiy remodellanishi hamda sistolik va diastolik disfunktsiyasi patogenezigacha hissa qo'shishi mumkin. Bizning tadqiqotimizda ham DKMP bilan og'rikan bemorlarda SYuYe og'irlik darajasi bilan qon zardobidagi C-reaktiv oqsil va IL-6, O'NO- α kabi yallig'lanisholdi sitokinlari o'rtasida yaqqol musbat bog'liqlik borligi ko'rsatildi.

Olingan natijalar dilatatsion kardiomiopatiya bilan og'rikan bemorlarda yallig'lanisholdi sitokinlari (IL-6, O'NO- α) va CRO kasallik kechishining og'irlik darajasiga baxo berishda muhim ahamiyatga ega ekanligini, hamda kelgusida bunday guruh bemorlarda yurak yetishmovchiligini

davolashda qo'llanilayotgan optimal medikamentoz terapiya tarkibiga kolxitsin dori vositasini qo'shish istiqbolli ekanligini ko'rsatadi.

Mualliflarning hissalari

Konseptualizatsiya, Abdullaev T.A.; metodologiya, Rahmatov M.M.; dasturiy ta'minot, Rahmatov M.M.; tasdiqlash, Abdullaev T.A., rasmiy tahlil, Abdullaev T.A.; tadqiqot, Rahmatov M.M.; resurslar, Rahmatov M.M.; ma'lumotlarni kuratorlik qilish, Abdullaev T.A.; original matnni yozish, Rahmatov M.M.; yozish va tahrirlash, Rahmatov M.M., G'ulomov X.A.; vizualizatsiya, Rahmatov M.M., G'ulomov X.A.; rahbarlik, Abdullaev T.A.; loyiha boshqaruvi, Abdullaev T.A.; moliya jalb qilish, Abdullaev T.A.. Barcha mualliflar nashr qilingan qo'lyozma versiyasi bilan tanish va u bilan rozi.

Authors' contribution.

Conceptualization, T.A. Abdullaev; methodology, M.M. Rakhmatov; software, M.M. Rakhmatov; validation, T.A. Abdullaev; formal analysis, T.A. Abdullaev; investigation, M.M. Rakhmatov; resources, M.M. Rakhmatov; data curation, T.A. Abdullaev; writing—original draft preparation, M.M. Rakhmatov; writing—review and editing, M.M. Rakhmatov, Kh.A. G'ulomov; visualization, M.M. Rakhmatov, Kh.A. G'ulomov; supervision, T.A. Abdullaev; project administration, T.A. Abdullaev; funding acquisition, T.A. Abdullaev. All authors have read and agreed to the published version of the manuscript.

Moliyalashtirish

Ishga tashqi moliya ajratilmagan.

Funding source.

The study was not funded.

Etika tamoyillariga muvofiqlik

Tadqiqot Xelsinki Deklaratsiyasiga muvofiq o'tkazildi va Respublika ixtisoslashtirilgan kardiologiya ilmiy-amaliy tibbiyot markazi tomonidan tasdiqlangan.

Ethics approval.

The study was conducted in accordance with the Declaration of Helsinki and was approved by the Republican Specialized Scientific-Practical Medical Center of Cardiology.

Nashrga xabardor qilingan rozilik.

Barcha tadqiqot ishtirokchilaridan xabardor qilingan rozilik olindi.

Consent for publication.

Informed consent was obtained from all study participants.

Ma'lumotlar mavjudligi to'g'risidagi bayonot

Taqdim etilgan barcha ma'lumot manbaalari adabiyotlar royxatida korsiqlangan.

Data Availability Statement

All data sources used are listed in the references section.

Rahmatnomalar

Mualliflar ushbu ilmiy ishni amalga oshirishda yordam bergan Respublika ixtisoslashtirilgan kardiologiya ilmiy-amaliy tibbiyot markazining Yurak yetishmovchiligi va miokard nokoronarogen kasalliklari laboratoriyasi xodimlariga chuqur minnatdorchilik bildiradilar.

Acknowledgments

The authors express their sincere gratitude to the staff of the Laboratory of Heart Failure and Non-Coronary Myocardial Diseases at the Republican Specialized Scientific and Practical Medical Center of Cardiology for their valuable assistance in carrying out this research work.

Manfaatlar to'qnashuvi

Mualliflar o'zlarining manfaatlar to'qnashuvi yo'qligini e'lon qiladilar.

Conflict of interest

The authors declare that they have no conflict of interest

Qisqartmalar

DKMP	Dilatatsion kardiomiopatiya
GKMP	Gipertrofik kardiomiopatiya
RKMP	Restriktiv kardiomiopatiya
AO'QKMP	Aritmogen o'ng qorincha kardiomiopatiyasi
ChQNDKMP	Chap qorincha nodilatatsion kardiomiopatiyasi
SYuYe	Surunkali yurak yetishmovchiligi
ChQ	Chap qorincha
OF	Otish fraksiyasi
CRO	C-reaktiv oqsil
IL-6	Interleykin-6
O'NO- α	O'sma nekrozi omili- α
NUP	Natriyuretik peptid
YuIK	Yurak ishemik kasalligi

Adabiyot

- [1] Arbelo E, Protonotarios A, Gimeno JR, Arbustini E, Barriales-Villa R, Basso C et al. ESC Scientific Document Group. 2023 ESC guidelines for the management of cardiomyopathies. *Eur Heart J* 2023; 44: 3503–3626.
- [2] Elliott P, Andersson B, Arbustini E. Classification of the cardiomyopathies: a position statement from the European Society of Cardiology working group on myocardial and pericardial diseases. *Eur Heart J* 2008; 29: 270–276.
- [3] Charron P, Arad M, Arbustini E, Basso C, Bilinska Z, Elliott P et al. ; European Society of Cardiology Working Group on Myocardial and Pericardial Diseases. Genetic counselling and testing in cardiomyopathies: a position statement of the European Society of Cardiology working group on myocardial and pericardial diseases. *Eur Heart J* 2010; 31: 2715–2726.
- [4] Rapezzi C, Arbustini E, Caforio AL, Charron P, Gimeno-Blanes J, Helio T et al. Diagnostic work-up in cardiomyopathies: bridging the gap between clinical phenotypes and final diagnosis. A position statement from the ESC working group on myocardial and pericardial diseases. *Eur Heart J* 2013; 34: 1448–1458.
- [5] Authors/Task Force members, Elliott PM, Anastakis A, Borger MA, Borggrefe M, Cecchi F, Charron P et al. 2014 ESC guidelines on diagnosis and management of hypertrophic cardiomyopathy: the task force for the diagnosis and management of hypertrophic cardiomyopathy of the European Society of Cardiology (ESC). *Eur Heart J* 2014; 35: 2733–2779.
- [6] Pinto YM, Elliott PM, Arbustini E, Adler Y, Anastakis A, Böhm et al. Proposal for a revised definition of dilated cardiomyopathy, hypokinetic non-dilated cardiomyopathy, and its implications for clinical practice: a position statement of the ESC working group on myocardial and pericardial diseases. *Eur Heart J* 2016; 37: 1850–1858.
- [7] Garcia-Pavia P, Rapezzi C, Adler Y, Arad M, Basso C, Brucato A et al. Diagnosis and treatment of cardiac amyloidosis: a position statement of the ESC working group on myocardial and pericardial diseases. *Eur Heart J* 2021; 42: 1554–1568.
- [8] Arbustini E, Behr ER, Carrier L, van Duijn C, Evans P, Favalli V et al. Interpretation and actionability of genetic variants in cardiomyopathies: a position statement from the European Society of Cardiology Council on cardiovascular genomics. *Eur Heart J* 2022; 43: 1901–1916.
- [9] Weintraub RG, Semsarian C, Macdonald P. Dilated cardiomyopathy. *Lancet*. 2017;390(10092):400–414. [https://doi.org/10.1016/S0140-6736\(16\)31713-5](https://doi.org/10.1016/S0140-6736(16)31713-5).
- [10] Elliott P, Andersson B, Arbustini E, Bilinska Z, Cecchi F, Charron P, et al. Classification of the cardiomyopathies: a position statement from the European Society of Cardiology Working Group on Myocardial and Pericardial Diseases. *Eur Heart J* 2008; 29: 270–276.
- [11] Seferovic PM, Polovina M, Bauersachs J, Arad M, Ben Gal T, Lund LH, Felix SB. et al. Heart failure in cardiomyopathies: a position paper from the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail*. 2019;21(5):553–576. <https://doi.org/10.1002/ejhf.1461>.
- [12] AHA/ACC/HFSA Guideline for the Management of Heart Failure. *Circulation*. 2022;145(18):e895–e1032.
- [13] 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. Th.McDonagh, M. Metra. <https://doi.org/10.15829/1560-4071-2023-5168>.
- [14] Belenkov Yu.N., Nasonov E.L., Samsonov M.Yu., Ageev F.T., Mareev V.Yu. // Heart Failure. 2000. Vol. 4. pp. 135–138.
- [15] Dinarello C.A. // Cytokine Growth Factor Reviews. 1997. V. 8 (4). P. 253–265.

- [16] Dick SA, Epelman S. Chronic heart failure and inflammation: what do we really know? *Circ Res.* 2016;119:159–176. <https://doi.org/10.1161/CIRCRESAHA.116.308030>.
- [17] Trachtenberg BH, Hare JM. Inflammatory cardiomyopathic syndromes. *Circ Res.* 2017;121:803–818. <https://doi.org/10.1161/CIRCRESAHA.117.310221>.
- [18] Wilson HM, Cheyne L, Brown PAJ, Kerr K, Hannah A, Srinivasan J, Duniak N, Horgan G, Dawson DK. Characterization of the myocardial inflammatory response in acute stress-induced (Takotsubo) cardiomyopathy. *JACC Basic Transl Sci.* 2018;3:766–778. <https://doi.org/10.1016/j.jacbts.2018.08.006>.
- [19] Shao Y., Redfors B., Scharin Tang M., Mollmann H., Troidl C., Szardien S. et al. Novel rat model reveals important roles of -adrenoreceptors in stress-induced cardiomyopathy. *International Journal of Cardiology.* 2013; 168(3):1943-50. <https://doi.org/10.1016/j.ijcard.2012.02.092>.
- [20] Sachdeva J., Dai W., Kloner R.A. Functional and Histological Assessment of an Experimental Model of Takotsubo's Cardiomyopathy. *Journal of American Heart Association.* 2014;3(3):e000921. <https://doi.org/10.1161/JAHA.114.000921>.
- [21] Ishikura F., Takano Y., Ueyama T. Acute effects of beta-blocker with intrinsic sympathomimetic activity on stress-induced cardiac dysfunction in rats. *Journal of Cardiology.* 2012;60(6): 470-4. <https://doi.org/10.1016/j.jcc.2012.07.004>.
- [22] Haggendal J., Johansson G., Jonsson L., Thoren-Tolling K. Effect of Propranolol on Myocardial Cell Necroses and Blood Levels of Catecholamines in Pigs Subjected to Stress. *Acta Pharmacologica et Toxicologica.* 2009;50(1): 58-66. <https://doi.org/10.1111/j.1600-0773.1982.tb00940.x>.
- [23] Kühl U, Noutsias M, Seeberg B, Schultheiss HP. Immunohistological evidence for a chronic intramyocardial inflammatory process in dilated cardiomyopathy. *Heart.* 1996;75:295–300.
- [24] Torre-Amione G, Kapadia S, Lee J, Durand JB, Bies RD, Young JB, Mann DL. Tumor necrosis factor- α and tumor necrosis factor receptors in the failing human heart. *Circulation.* 1996;93:704–711. <https://doi.org/10.1161/01.cir.93.4.704>.
- [25] Belenkov Yu.N., Ageev F.T., Mareev V.Yu. Neurohormones and cytokines in heart failure: a new theory of an old disease? *Journal Heart Failure*, 2000: 1(4), 35–38.
- [26] Dunlay SM, Weston SA, Redfield MM, Killian JM, Roger VL. Tumor necrosis factor- α and mortality in heart failure: a community study. *Circulation.* 2008;118:625–631. <https://doi.org/10.1161/CIRCULATIONAHA.107.759191>.
- [27] Dewald O, Ren G, Duerr GD, Zoerlein M, Klemm C, Gersch C, Tincey S, Michael LH, Entman ML, Frangogiannis NG. Of mice and dogs: species-specific differences in the inflammatory response following myocardial infarction. *Am J Pathol.* 2004;164:665–677. [https://doi.org/10.1016/S0002-9440\(10\)63154-9](https://doi.org/10.1016/S0002-9440(10)63154-9).
- [28] Shioi T, Matsumori A, Kihara Y, Inoko M, Ono K, Iwanaga Y, Yamada T, Iwasaki A, Matsushima K, Sasayama S. Increased expression of interleukin-1 β and monocyte chemoattractant and activating factor/monocyte chemoattractant protein-1 in the hypertrophied and failing heart with pressure overload. *Circ Res.* 1997;81:664–671. <https://doi.org/10.1161/01.res.81.5.664>.
- [29] Francis SE, Holden H, Holt CM, Duff GW. Interleukin-1 in myocardium and coronary arteries of patients with dilated cardiomyopathy. *J Mol Cell Cardiol.* 1998;30:215–223. <https://doi.org/10.1006/jmcc.1997.0592>.
- [30] Shioi T, Matsumori A, Kihara Y, Inoko M, Ono K, Iwanaga Y, Yamada T, Iwasaki A, Matsushima K, Sasayama S. Increased expression of interleukin-1 β and monocyte chemoattractant and activating factor/monocyte chemoattractant protein-1 in the hypertrophied and failing heart with pressure overload. *Circ Res.* 1997;81:664–671. <https://doi.org/10.1161/01.res.81.5.664>.
- [31] Suetomi T, Willeford A, Brand CS, Cho Y, Ross RS, Miyamoto S, Brown JH. Inflammation and NLRP3 inflammasome activation initiated in response to pressure overload by Ca(2+)/calmodulin-dependent protein kinase II δ signaling in cardiomyocytes are essential for adverse cardiac remodeling. *Circulation.* 2018;138:2530–2544. <https://doi.org/10.1161/CIRCULATIONAHA.118.034621>.
- [32] Kawaguchi M, Takahashi M, Hata T, Kashima Y, Usui F, Morimoto H, Izawa A, Takahashi Y, Masumoto J, Koyama J, Hongo M, Noda T, Nakayama J, Sagara J, Taniguchi S, Ikeda U. Inflammasome activation of cardiac fibroblasts is essential for myocardial ischemia/reperfusion injury. *Circulation.* 2011;123:594–604. <https://doi.org/10.1161/CIRCULATIONAHA.110.982777>.
- [33] Murphy SP, Kakkar R, McCarthy CP, Januzzi JL., Jr Inflammation in heart failure: JACC state-of-the-art review. *J Am Coll Cardiol.* 2020;75(11):1324–40. <https://doi.org/10.1016/j.jacc.2020.01.014>.
- [34] Liu S.J., Zhou W., Kennedy R.H. Suppression of beta-adrenergic responsiveness of L-type Ca²⁺ current by IL-1 β in rat ventricular myocytes. *Am J Physiol.* 1999; 276(1):H141-8. <https://doi.org/10.1152/ajpheart.1999.276.1.H141>.
- [35] Tatsumi T., Matoba S., Kawahara A. et al. Cytokine-induced nitric oxide production inhibits mitochondrial energy production and impairs contractile function in rat cardiac myocytes. *Pflugers Arch.* 2002; 35(5): 1338-46. [https://doi.org/10.1016/s0735-1097\(00\)00526-x](https://doi.org/10.1016/s0735-1097(00)00526-x).

- [36] Aleksova A., Beltrami A.P., Carriere C. et al. Interleukin-1 levels predict long-term mortality and need for heart transplantation in ambulatory patients affected by idiopathic dilated cardiomyopathy. *Oncotarget*. 2017; 8(15):25131-25140. <https://doi.org/10.18632/oncotarget.15349>.
- [37] Kubota T, Miyagishima M, Alvarez RJ, et al. Expression of proinflammatory cytokines in the failing human heart: comparison of recent-onset and end-stage congestive heart failure. *J Heart Lung Transplant*. 2000;19:819–824. [https://doi.org/10.1016/s1053-2498\(00\)00173-x](https://doi.org/10.1016/s1053-2498(00)00173-x).
- [38] Raymond RJ, Dehmer GJ, Theoharides TC, Deliargyris EN. Elevated interleukin-6 levels in patients with asymptomatic left ventricular systolic dysfunction. *Am Heart J*. 2001;141(3):435–8. <https://doi.org/10.1067/mhj.2001.113078>.
- [39] Vasan RS, Sullivan LM, Roubenoff R, Dinarello CA, Harris T, Benjamin EJ, et al. Inflammatory markers and risk of heart failure in elderly subjects without prior myocardial infarction: the Framingham Heart Study. *Circulation*. 2003;107(11):1486–91. <https://doi.org/10.1161/01.CIR.0000057810.48709.F6>.
- [40] Yamauchi-Takahara K, Ihara Y, Ogata A, Yoshizaki K, Azuma J, Kishimoto T. Hypoxic stress induces cardiac myocyte–derived interleukin-6. *Circulation*. 1995;91(5):1520–4. <https://doi.org/10.1161/01.CIR.91.5.1520>.
- [41] Pfisterer M, Buser P, Rickli H, Gutmann M, Erne P, Rickenbacher P, et al. BNP-guided vs symptom-guided heart failure therapy: the trial of intensified vs Standard Medical Therapy in Elderly patients with congestive heart failure (TIME-CHF) randomized trial. *JAMA*. 2009;301(4):383–92. <https://doi.org/10.1001/jama.2009.2>.
- [42] Imazio M, Nidorf M. Colchicine and the heart. *Eur Heart J*. 2021;42(28):2745-2760.
- [43] Deftereos C, Giannopoulos G, Panagopoulos V, Bouras G, Raisakis R, Koskinas K, et al. Anti-Inflammatory Treatment With Colchicine in Stable Chronic Heart Failure. *JACC Heart Fail*. 2014;2(2):131-137.
- [44] Pascual-Figal D, Núñez J, Pérez-Martínez MT, González-Juanatey JR, Taibo-Urquía M, Llacer-Borrà P, et al. Colchicine in acutely decompensated heart failure: the COLICA trial. *Eur Heart J*. 2024;45(45):4826-4836.
- [45] Xuan Sun et al., Colchicine Ameliorates Dilated Cardiomyopathy Via SIRT2-Mediated Suppression of NLRP3 Inflammasome Activation. *JAHA*, Volume 11, Number 13, <https://doi.org/10.1161/JAHA.122.025266>.

Nashriyot javobgar emas/ eslatmasi:

Barcha nashrlarda keltirilgan bayonotlar, fikrlar va ma'lumotlar faqat mualliflar va ishtirokchilarga tegishlidir, na Jurnal va na muharrirlar. Jurnal va muharrirlar, mazkur kontentda keltirilgan har qanday g'oyalar, usullar, ko'rsatmalar yoki mahsulotlar natijasida insonlar yoki mulkka yetkazilgan har qanday zarar uchun javobgar emas.

Disclaimer of liability/Publisher's Note: The statements, opinions and data contained in all publications belong exclusively to individuals. The authors and participants, and the Journal and the editors. The journal and the editors are not responsible for any damage caused to people or property resulting from any ideas, methods, instructions or products mentioned in the content.